

SERCA2 Rabbit mAb

货号: **AYM28519**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	109kDa/111kDa/114kDa
实测分子量	110,140kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,NIH/3T3,Mouse heart,Rat heart
细胞定位	Endoplasmic reticulum membrane,Multi-pass membrane protein,Sarcoplasmic reticulum membrane
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human SERCA2.
序列	MENAHTKTVEEVLGHFGVNESTGLSLEQVKLKERWGSNELPAEEGKTLLELVIEQFEDLLVRILLLAACISFVLAWFEE

靶点信息

研究背景	This gene encodes one of the SERCA Ca(2+)-ATPases, which are intracellular pumps located in the sarcoplasmic or endoplasmic reticula of muscle cells. This enzyme catalyzes the hydrolysis of ATP coupled with the translocation of calcium from the cytosol into the sarcoplasmic reticulum lumen, and is involved in regulation of the contraction/relaxation cycle. Mutations in this gene cause Darier-White disease, also known as keratosis follicularis, an autosomal dominant skin disorder characterized by loss of adhesion between epidermal cells and abnormal keratinization. Alternative splicing results in multiple transcript variants encoding different isoforms.
基因ID	488
基因名	ATP2A2
Swiss	P16615
别名	ATP2B;DAR;DD;SERCA2;ATP2A2/SERCA2;ATP2A2

产品验证



Western blot analysis of SERCA2 expressed in Mouse heart,Rat heart,293T using SERCA2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn