

# Cullin4A Rabbit mAb

货号: **AYM28525**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC IP FC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>ICC:</b> 1:50 - 1:200 <b>IP:</b> 1:20 - 1:50 <b>FC:</b> 1:20 - 1:50
理论分子量	76kDa/87kDa
实测分子量	88kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562,BT-474,HeLa,Mouse testis
细胞定位	nucleoplasm,nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Cullin4A.
序列	IERERSGEAVDRSLLRSLLGMLSDLQVYKDSFELKFLEETNCLYAAEQRLMQEREVPEYLNHVSKRLEEEGDRVITYLDHS TQKPLIACVEKQLLGEHLT

## 靶点信息

研究背景	CUL4A is the ubiquitin ligase component of a multimeric complex involved in the degradation of DNA damage-response proteins (Liu et al., 2009 [PubMed 19481525]).[supplied by OMIM, Oct 2009]
基因ID	8451

基因名	CUL4A
Swiss	Q13619
别名	CUL4A;cullin-4A

## 产品验证



Western blot analysis of Cullin4A expressed in K-562, BT-474, HeLa using Cullin4A Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)