— ABLYBIO, Help Your Research

CD97 Rabbit mAb

货号: AYM28542

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	**WB:** 1:500 - 1:2000 **IHC:** 1:50 - 1:200 **IF/ICC:** 1:50 - 1:200 **IP:** 1:20 - 1:50 **FC:** 1:20 - 1:50
理论分子量	81kDa/86kDa/91kDa
实测分子量	92kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,Jurkat,Mouse lung,Rat liver
细胞定位	Cell membrane,Multi-pass membrane protein,Secreted,extracellular space
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein corresponding to Human CD97.

靶点信息

研究背景	This gene encodes a member of the EGF-TM7 subfamily of adhesion G protein-coupled receptors, which mediate cell-cell interactions. These proteins are cleaved by self-catalytic proteolysis into a large extracel lular subunit and seven-span transmembrane subunit, which associate at the cell surface as a receptor c omplex. The encoded protein may play a role in cell adhesion as well as leukocyte recruitment, activation and migration, and contains multiple extracellular EGF-like repeats which mediate binding to chondroitin sulfate and the cell surface complement regulatory protein CD55. Expression of this gene may play a role in the progression of several types of cancer. Alternatively spliced transcript variants encoding multiple is oforms with 3 to 5 EGF-like repeats have been observed for this gene. This gene is found in a cluster with other EGF-TM7 genes on the short arm of chromosome 19.
基因ID	976
基因名	CD97
Swiss	P48960 (https://www.uniprot.org/uniprotkb/P48960/entry)
别名	CD97,CD97 Rabbit mAb,Leukocyte antigen CD97

产品验证



Western blot analysis of CD97 expressed in Mouse lung,HepG2,Jurkat using CD97 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)