

IKK alpha Rabbit mAb

货号: **AYM28549**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	84kDa
实测分子量	85kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,Rat kidney
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human IKK alpha.
序列	VLKELFGHLSKLLGCKQKIIDLDPKVEVALSNIKEADNTVMFMQGKRQKEIWHLLKIACTQSSARSLVGSSEGAVTPQ TSA WLPPTSAEHDHSLSCVVTP

靶点信息


研究背景	This gene encodes a member of the serine/threonine protein kinase family. The encoded protein, a component of a cytokine-activated protein complex that is an inhibitor of the essential transcription factor NF-kappa-B complex, phosphorylates sites that trigger the degradation of the inhibitor via the ubiquitination pathway, thereby activating the transcription factor.
------	---

基因ID	1147
基因名	CHUK
Swiss	O15111
别名	CHUK;IKBKA;IKK-alpha;IKK1;IKKA;NFKBIKA;TCF16;IKK alpha

产品验证



Immunohistochemical analysis of paraffin-embedded mouse kidney, using IKK alpha Antibody.



Western blot analysis of IKK alpha expressed in 293T,Rat kidney using IKK alpha Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn