

— ABLYBIO, Help Your Research



# ASK1 Rabbit mAb

货号: **AYM28570**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200 <b>FC:</b> 1:20 - 1:50
理论分子量	69kDa/154kDa
实测分子量	155kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-431,HeLa
细胞定位	Cytoplasm,Endoplasmic reticulum
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human ASK1 .
------	--


## 靶点信息

研究背景	Mitogen-activated protein kinase (MAPK) signaling cascades include MAPK or extracellular signal-regulated kinase (ERK), MAPK kinase (MKK or MEK), and MAPK kinase kinase (MAPKKK or MEKK). MAPKK kinase/MEKK phosphorylates and activates its downstream protein kinase, MAPK kinase/MEK, which in turn activates MAPK. The kinases of these signaling cascades are highly conserved, and homologs exist in yeast, <i>Drosophila</i> , and mammalian cells. MAPKKK5 contains 1,374 amino acids with all 11 kinase subdomains. Northern blot analysis shows that MAPKKK5 transcript is abundantly expressed in human heart and pancreas. The MAPKKK5 protein phosphorylates and activates MKK4 (aliases SERK1, MAPKK4) <i>in vitro</i> , and activates c-Jun N-terminal kinase (JNK)/stress-activated protein kinase (SAPK) during transient expression in COS and 293 cells; MAPKKK5 does not activate MAPK/ERK.
基因ID	4217
基因名	MAP3K5
Swiss	Q99683 ( <a href="https://www.uniprot.org/uniprotkb/Q99683/entry">https://www.uniprot.org/uniprotkb/Q99683/entry</a> )
别名	ASK1,ASK1 Rabbit mAb,MAP3K5,Apoptosis signal-regulating kinase 1,MAPK/ERK kinase kinase 5,MAPKKK5,MEKK5


## 产品验证




Western blot analysis of ASK1 expressed in A-431, HeLa using ASK1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.




Immunohistochemical analysis of paraffin-embedded Rat stomach, using the Antibody at 1:50 dilution.




Immunohistochemical analysis of paraffin-embedded Human Hodgkin's lymphoma, using the Antibody at 1:50 dilution.




Immunohistochemical analysis of paraffin-embedded Human prostate cancer, using the Antibody at 1:50 dilution.



Immunohistochemical analysis of paraffin-embedded Human astrocytoma, using the Antibody at 1:50 dilution.



Immunohistochemical analysis of paraffin-embedded Mouse skin, using the Antibody at 1:50 dilution.



Immunohistochemical analysis of paraffin-embedded human breast cancer, using ASK1 Antibody.

### 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)