

— ABLYBIO, Help Your Research



NR0B1 / DAX1 Rabbit mAb

货号: **AYM28576**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	43kDa/51kDa
实测分子量	52kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,NCI-H460,MCF7
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human NR0B1 / DAX1.
------	---

靶点信息

研究背景	This gene encodes a protein that contains a DNA-binding domain. The encoded protein acts as a dominant-negative regulator of transcription which is mediated by the retinoic acid receptor. This protein also functions as an anti-testis gene by acting antagonistically to Sry. Mutations in this gene result in both X-linked congenital adrenal hypoplasia and hypogonadotropic hypogonadism.
基因ID	190
基因名	NR0B1
Swiss	P51843 (https://www.uniprot.org/uniprotkb/P51843/entry)
别名	NR0B1 / DAX1, NR0B1 / DAX1 Rabbit mAb, NR0B1, DSS-AHC critical region on the X chromosome protein 1, Nuclear receptor DAX-1, AHC, DAX1

产品验证



Western blot analysis of NR0B1 / DAX1 expressed in U-87MG, NCI-H460, MCF7 using NR0B1 / DAX1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)