

— ABLYBIO, Help Your Research



HMGCL Rabbit mAb

货号: **AYM28577**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	20kDa/26kDa/34kDa
实测分子量	30kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,SKOV3,BT-474,MCF7,Mouse liver,Mouse kidney
细胞定位	Mitochondrion matrix,Peroxisome
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human HMGCL.
------	----------------------------------------------------------

靶点信息

研究背景	The protein encoded by this gene belongs to the HMG-CoA lyase family. It is a mitochondrial enzyme that catalyzes the final step of leucine degradation and plays a key role in ketone body formation. Mutations in this gene are associated with HMG-CoA lyase deficiency. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
基因ID	3155
基因名	HMGCL
Swiss	P35914 (https://www.uniprot.org/uniprotkb/P35914/entry)
别名	HMGCL,HMGCL Rabbit mAb,3-hydroxy-3-methylglutarate-CoA lyase

产品验证



Western blot analysis of HMGCL expressed in U-251MG,SKOV3,BT-474 using HMGCL Rabbit mAb at 1:100. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)