

CDX2 Rabbit mAb

货号: **AYM28579**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	33kDa
实测分子量	40kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SW620,MCF7
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human CDX2.
序列	MYVSYLLDKDVSMYPSSVRHSGGLNLPQNFVSPQYDPDYGGYHVAAAAAAAAANLDSAQSPGPSWPAAYGAPLREDW NGYAPGGAAAAANAVAHGLNGGSPAAMGYSSPADYHPHHHPHHHPHAAAPSCASGLLQTLNPGPPGAATAAAEQ LSPGGQRRNLCEWMRKPAQQLGSQ

靶点信息


研究背景	This gene is a member of the caudal-related homeobox transcription factor gene family. The encoded protein is a major regulator of intestine-specific genes involved in cell growth and differentiation. This protein also plays a role in early embryonic development of the intestinal tract. Aberrant expression of this gene is associated with intestinal inflammation and tumorigenesis.
------	--

基因ID	1045
基因名	CDX2
Swiss	Q99626
别名	CDX2;CDX-3;CDX2/AS;CDX3

产品验证



Immunohistochemical analysis of paraffin-embedded human colon, using CDX2 Antibody.



Western blot analysis of CDX2 expressed in SW620, MCF7 using CDX2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn