

— ABLYBIO, Help Your Research



# Bcl10 Rabbit mAb

货号: **AYM28586**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200 <b>IP:</b> 1:20 - 1:50
理论分子量	26kDa
实测分子量	31kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji,Mouse spleen,Rat spleen
细胞定位	Cytoplasm,Membrane raft,perinuclear region
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Bcl10.
------	--


## 靶点信息

研究背景	This gene was identified by its translocation in a case of mucosa-associated lymphoid tissue (MALT) lymphoma. The protein encoded by this gene contains a caspase recruitment domain (CARD), and has been shown to induce apoptosis and to activate NF-kappaB. This protein is reported to interact with other CARD domain containing proteins including CARD9, 10, 11 and 14, which are thought to function as upstream regulators in NF-kappaB signaling. This protein is found to form a complex with MALT1, a protein encoded by another gene known to be translocated in MALT lymphoma. MALT1 and this protein are thought to synergize in the activation of NF-kappaB, and the deregulation of either of them may contribute to the same pathogenetic process that leads to the malignancy. Alternative splicing results in multiple transcript variants.
基因ID	8915
基因名	BCL10
Swiss	O95999 ( <a href="https://www.uniprot.org/uniprotkb/O95999/entry">https://www.uniprot.org/uniprotkb/O95999/entry</a> )
别名	Bcl10,Bcl10 Rabbit mAb,B-cell CLL/lymphoma 10,CARD-containing molecule enhancing NF-kappa-B,CARD-like apoptotic protein,CED-3/ICH-1 prodomain homologous E10-like regulator,Cellular homolog of vCARMEN,Cellular-E10

## 产品验证



Western blot analysis of Bcl10 expressed in Raji, Mouse spleen, Rat spleen using Bcl10 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



Immunohistochemical analysis of paraffin-embedded human breast cancer, using Bcl10 Antibody.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))