

— ABLYBIO, Help Your Research



DDX58 Rabbit mAb

货号: **AYM28598**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IP
推荐浓度	WB: 1:500 - 1:2000 IP: 1:20 - 1:50
理论分子量	101kDa/106kDa
实测分子量	107kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2
细胞定位	Cell junction,Cell projection,Cytoplasm,cytoskeleton,ruffle membrane,tight junction
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human DDX58.
------	--

靶点信息

研究背景	DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases which are implicated in a number of cellular processes involving RNA binding and alteration of RNA secondary structure. This gene encodes a protein containing RNA helicase-DEAD box protein motifs and a caspase recruitment domain (CARD). It is involved in viral double-stranded (ds) RNA recognition and the regulation of immune response.
基因ID	23586
基因名	DDX58
Swiss	O95786 (https://www.uniprot.org/uniprotkb/O95786/entry)
别名	DDX58, DDX58 Rabbit mAb, ATP-dependent RNA helicase DDX58, DEAD box protein 58, RIG-I-like receptor 1, RNA sensor RIG-I, Retinoic acid-inducible gene 1 protein, Retinoic acid-inducible gene I protein

产品验证



Western blot analysis of DDX58 expressed in HepG2 using DDX58 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)