

Methyl-Histone H3 (di K4) Rabbit mAb

货号: **AYM28619**

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse,Mouse,Rat,Rat |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IHC IF/ICC IP FC CHIP |
| 推荐浓度 | WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50 |
| 理论分子量 | 15kDa |
| 实测分子量 | 15kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HeLa |
| 细胞定位 | Chromosome,Nucleus |
| 纯化 | Affinity purification |


抗原信息

| | |
|------|--|
| 抗原信息 | Recombinant fusion protein corresponding to Human Methyl-Histone H3 (di K4). |
| 序列 | PRKQL |


靶点信息

| | |
|-------|---|
| 研究背景 | Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the small histone gene cluster on chromosome 6p22-p21.3. |
| 基因ID | 8356 |
| 基因名 | |
| Swiss | P68431 |
| 别名 | HIST1H3J;H3/j;H3FJ |

产品验证



Immunohistochemical analysis of paraffin-embedded human liver, using Methyl-Histone H3 (di K4) Antibody.



Western blot analysis of Methyl-Histone H3 (di K4) expressed in HeLa using Methyl-Histone H3 (di K4) Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn