

CHD4 Rabbit mAb

货号: **AYM28655**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	218kDa/220kDa
实测分子量	260kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	HeLa, Jurkat, 293T, HT-1080
细胞定位	Cytoplasm, Nucleus, centrosome, cytoskeleton, microtubule organizing center
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human CHD4.
序列	ELAEVEENKKMSQGPSKPTPTSTPGDTPNTAPVPPAEDGKIEENSLKEEESIEGEKEVKSTAPETAIECTQAPAPASE DEKVVVEPPEGEEKVEKAEVKERTEEPMETEPKGAADVEKVEEKSAIDLTPIVVEDKEEKKEEEEKVEMLQNGETPKDLND EKQKK

靶点信息

研究背景	The product of this gene belongs to the SNF2/RAD54 helicase family. It represents the main component of the nucleosome remodeling and deacetylase complex and plays an important role in epigenetic transcriptional repression. Patients with dermatomyositis develop antibodies against this protein. Somatic mutations in this gene are associated with serous endometrial tumors. Alternative splicing results in multiple transcript variants encoding different isoforms.
------	--

基因ID	1108
基因名	CHD4
Swiss	Q14839
别名	CHD4;CHD-4;Mi-2b;Mi2-BETA;SIHIWES

产品验证



Western blot analysis of CHD4 expressed in HeLa, Jurkat, 293T using CHD4 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn