

STK33 Rabbit mAb

货号: **AYM28657**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	58, 50kDa
实测分子量	58, 50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y
细胞定位	cytoplasm,nucleus,perinuclear region of cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human STK33.
序列	KEWKNNPESVEENTTEEKKNKPSTEEKLKSYPWGNVPDANYTSDEEEEKQSTAYEKQFPATSKDNFDMCSSFTSSKLLP AEIKGEMEKTPVTPSQGTATKYPKSGALSRTKKKL

靶点信息

研究背景	Serine/threonine protein kinase which phosphorylates VIME. May play a specific role in the dynamic behavior of the intermediate filament cytoskeleton by phosphorylation of VIME (By similarity. Not essential for the survival of KRAS-dependent AML cell lines.
基因ID	65975

基因名	STK33
Swiss	Q9BYT3
别名	

产品验证



Western blot analysis of STK33 expressed in SH-SY5Y using STK33 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn