— ABLYBIO, Help Your Research

DDAH1 Rabbit mAb

货号: AYM28682

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	**WB:** 1:500 - 1:2000 **IHC:** 1:50 - 1:200 **IF/ICC:** 1:50 - 1:200 **IP:** 1:20 - 1:50 **FC:** 1:20 - 1:50
理论分子量	20kDa/31kDa
实测分子量	37kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SW480,22Rv1,HT-29,293T,SGC-7901,Mouse brain,Mouse kidney,Rat brain
细胞定位	cytosol,extracellular exosome
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human DDAH1.

靶点信息

研究背景	This gene belongs to the dimethylarginine dimethylaminohydrolase (DDAH) gene family. The encoded en zyme plays a role in nitric oxide generation by regulating cellular concentrations of methylarginines, whic h in turn inhibit nitric oxide synthase activity.
基因ID	23576
基因名	DDAH1
Swiss	O94760 (https://www.uniprot.org/uniprotkb/O94760/entry)
别名	DDAH1,DDAH1 Rabbit mAb,DDAHI,Dimethylargininase-1,DDAH

产品验证



Western blot analysis of DDAH1 expressed in SW480,22Rv1,HT-29 using DDAH1 Rabbit mAb at 1:1000. Se condary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking bu ffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)