

DDAH1 Rabbit mAb

货号: **AYM28682**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	20kDa/31kDa
实测分子量	37kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SW480,22Rv1,HT-29,293T,SGC-7901,Mouse brain,Mouse kidney,Rat brain
细胞定位	cytosol,extracellular exosome
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human DDAH1.
序列	MAGLGHPAAFGRATHAVVRALPESLGQHALRSAKGEEVDVARAERQHQLYVGVGLGSKLGLQVVLPADSLPDCVFVED VAVVCEETALITRPGAPSRRKEVDMMEALEKLQNLIVEMKDENATLDGGDVLFTGREFFVGLSKRTNQRGAELADTFKD YAVSTVPVADGLHLKSFCSMAGPNLIAIGSSESAQKALKIMQQMSDHRDYDKLTPDDIAANCIYLNIPNKGHVLLHRTPEEY PESAKVYEKLDHMLIPVSMSELEKVDGLLTCCSVLINKKVDS

靶点信息

研究背景	This gene belongs to the dimethylarginine dimethylaminohydrolase (DDAH) gene family. The encoded enzyme plays a role in nitric oxide generation by regulating cellular concentrations of methylarginines, which in turn inhibit nitric oxide synthase activity.
------	---

基因ID	23576
基因名	DDAH1
Swiss	O94760
别名	DDAH1;DDAH;DDAH-1;DDAHI;HEL-S-16

产品验证



Western blot analysis of DDAH1 expressed in SW480,22Rv1,HT-29 using DDAH1 Rabbit mAb at 1:1000 . Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blo cking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn