

# DDAH2 Rabbit mAb

货号: **AYM28686**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IP
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IP:</b> 1:20 - 1:50
理论分子量	29kDa
实测分子量	30kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SW620,BT-474,MCF7,Mouse lung,Rat brain
细胞定位	Cytoplasm,Mitochondrion
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human DDAH2.
序列	MGTPGEGLGRCSHALIRGVPESLASGEGAGAGLPALDLAKAQREHGVGGKLRQRLGLQLLELPPEESLPLGPLLGD TAVI QGDTALITRPWSPARRPEVDGVRKALQDLGLRIVEIGDENATLDGTDVLF TGREFFVGLSKWTNHRGAEIVADTFRDFAV STVPVSGPSHLRGLCGMGGPRTVVAGSSDAAQKAVRAMAVLTDHPYASLTLPDDAAADCLFLRPGLPGVPPFLLHRGGG DLPNSQEALQKLSDVTLVPVSCSELEKAGAGLSSLCLVLSTRPHS

## 靶点信息

研究背景	This gene encodes a dimethylarginine dimethylaminohydrolase. The encoded enzyme functions in nitric oxide generation by regulating the cellular concentrations of methylarginines, which in turn inhibit nitric oxide synthase activity. The protein may be localized to the mitochondria. Alternative splicing resulting in multiple transcript variants.
------	--

基因ID	23564
基因名	DDAH2
Swiss	O95865
别名	DDAH2;DDAH;DDAHII;G6a;HEL-S-277;NG30

## 产品验证



Western blot analysis of DDAH2 expressed in Mouse lung,Rat brain,SW620 using DDAH2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)