

— ABLYBIO, Help Your Research



ITCH Rabbit mAb

货号: **AYM28688**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	103kDa
实测分子量	103kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,293T,Raji,NIH/3T3,C6,Mouse lung
细胞定位	cell cortex,cytoplasm,cytoplasmic vesicle,cytosol,extracellular exosome,nucleoplasm,plasma membrane
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human ITCH .
------	--

靶点信息

研究背景	This gene encodes a member of the Nedd4 family of HECT domain E3 ubiquitin ligases. HECT domain E3 ubiquitin ligases transfer ubiquitin from E2 ubiquitin-conjugating enzymes to protein substrates, thus targeting specific proteins for lysosomal degradation. The encoded protein plays a role in multiple cellular processes including erythroid and lymphoid cell differentiation and the regulation of immune responses. Mutations in this gene are a cause of syndromic multisystem autoimmune disease. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Mar 20 12]
基因ID	83737
基因名	ITCH
Swiss	Q96J02 (https://www.uniprot.org/uniprotkb/Q96J02/entry)
别名	ITCH,ITCH Rabbit mAb,Atrophin-1-interacting protein 4,HECT-type E3 ubiquitin transferase Itchy homolog, NFE2-associated polypeptide 1

产品验证



Western blot analysis of ITCH expressed in Mouse lung, HeLa, 293T using ITCH Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)