

DCTN5 Rabbit mAb

货号: **AYM28705**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	9kDa/16kDa/20kDa
实测分子量	22kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,293T,Mouse eye,Mouse testis,Mouse kidney,Rat lung,Rat spleen
细胞定位	Chromosome,Cytoplasm,centromere,cytoskeleton,kinetochore
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human DCTN5.
序列	MELGELLYNKSEYIETASGNKVSRSQSVLCGSQNIIVLNGKTIVMNDCCIIRGDLANVRVGRHCVVKRSRVIRPPFKKFSKGVAF FPLHIGDHFIEEDCVVN

靶点信息

研究背景	This gene encodes a subunit of dynactin, a component of the cytoplasmic dynein motor machinery involved in minus-end-directed transport. The encoded protein is a component of the pointed-end subcomplex and is thought to bind membranous cargo. A pseudogene of this gene is located on the long arm of chromosome 1. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.
------	---

基因ID	84516
基因名	DCTN5
Swiss	Q9BTE1
别名	DCTN5

产品验证



Western blot analysis of DCTN5 expressed in Mouse eye,Rat lung,MCF7 using DCTN5 Rabbit mAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn