

# DCTN5 Rabbit mAb

货号: **AYM28705**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200 <b>IP:</b> 1:20 - 1:50
理论分子量	9kDa/16kDa/20kDa
实测分子量	22kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,293T,Mouse eye,Mouse testis,Mouse kidney,Rat lung,Rat spleen
细胞定位	Chromosome,Cytoplasm,centromere,cytoskeleton,kinetochore
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human DCTN5.
------	--

## 靶点信息

研究背景	This gene encodes a subunit of dynactin, a component of the cytoplasmic dynein motor machinery involved in minus-end-directed transport. The encoded protein is a component of the pointed-end subcomplex and is thought to bind membranous cargo. A pseudogene of this gene is located on the long arm of chromosome 1. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.
基因ID	84516

基因名	DCTN5
Swiss	Q9BTE1
别名	DCTN5

## 产品验证



Western blot analysis of DCTN5 expressed in Mouse eye,Rat lung,MCF7 using DCTN5 Rabbit mAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)