

— ABLYBIO, Help Your Research



PDLIM1 Rabbit mAb

货号: **AYM28708**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	36kDa
实测分子量	36kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BxPC-3,THP-1,Mouse thymus
细胞定位	Cytoplasm,cytoskeleton
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human PDLIM1.
------	---

靶点信息

研究背景	This gene encodes a member of the enigma protein family. The protein contains two protein interacting domains, a PDZ domain at the amino terminal end and one to three LIM domains at the carboxyl terminal. It is a cytoplasmic protein associated with the cytoskeleton. The protein may function as an adapter to bring other LIM-interacting proteins to the cytoskeleton. Pseudogenes associated with this gene are located on chromosomes 3, 14 and 17.
基因ID	9124
基因名	PDLIM1
Swiss	O00151 (https://www.uniprot.org/uniprotkb/O00151/entry)
别名	PDLIM1,PDLIM1 Rabbit mAb,C-terminal LIM domain protein 1,Elfin,LIM domain protein CLP-36,CLIM1,CLP36

产品验证



Western blot analysis of PDLIM1 expressed in BxPC-3, THP-1, Mouse thymus using PDLIM1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)