

PDCD4 Rabbit mAb

货号: **AYM28712**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	50kDa/51kDa
实测分子量	52kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,HeLa,MCF7,Mouse thymus
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human PDCD4.
序列	MDVENEQILNVNPADPDNLSDSLFSGDEENAGTEEIKNEINGNWISASSINEARINAKAKRRLRKNSRDSGRGDSVSDS GSDALRSGLTVPSPKGRLLDRRSRSGKGRGLPKGGAGGKGVWGTGPGQVYDVEEVDVKDPNYDDDQENCVYETVVL PLDERAFEKTLTPIIQEYFEHGDTNEVAEMLRDLNLGEMKSGVPVLAVSLALEGKASHREMTSKLLSDLCGTMSTTDVEK SFDKLLKDLPELALDTPRAPQL

靶点信息

研究背景	This gene is a tumor suppressor and encodes a protein that binds to the eukaryotic translation initiation factor 4A1 and inhibits its function by preventing RNA binding. Alternative splicing results in multiple transcript variants.
------	---

基因ID	27250
基因名	PDCD4
Swiss	Q53EL6
别名	PDCD4;H731

产品验证



Western blot analysis of PDCD4 expressed in Mouse thymus,293T,HeLa using PDCD4 Rabbit mAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. B locking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn