

CD25 Rabbit mAb

货号: **AYM28714**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	IHC IF/ICC
推荐浓度	IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	30kDa
实测分子量	60KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat spleen,Rat lung
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human CD25.
序列	EQKERKTTEMQSPMQPVDQASLPGHCREPPPWENEATERIYHFVVGQMVYYQCVQGYRALHRGPAESVCKMTHGKTRW TQPQLICTGEMETSQFPGEEKPQ

靶点信息

研究背景	The interleukin 2 (IL2) receptor alpha (IL2RA) and beta (IL2RB) chains, together with the common gamma chain (IL2RG), constitute the high-affinity IL2 receptor. Homodimeric alpha chains (IL2RA) result in low-affinity receptor, while homodimeric beta (IL2RB) chains produce a medium-affinity receptor. Normally an integral-membrane protein, soluble IL2RA has been isolated and determined to result from extracellular proteolysis. Alternately-spliced IL2RA mRNAs have been isolated, but the significance of each is presently unknown. Mutations in this gene are associated with interleukin 2 receptor alpha deficiency.
------	---

基因ID	3559
基因名	IL2RA
Swiss	P01589
别名	IL2RA;CD25;IDDM10;IL2R;IMD41;TCGFR;p55

产品验证



Immunohistochemical analysis of paraffin-embedded human tonsil, using CD25 Antibody.

实验步骤

访问官网浏览详情: www.ablybio.cn