

CRABP2 Rabbit mAb

货号: **AYM28719**

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IHC IF/ICC FC |
| 推荐浓度 | WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50 |
| 理论分子量 | 15kDa |
| 实测分子量 | 14kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | MCF7,SW480,A-431,22Rv1,HT-29,293T,Mouse brain,Mouse uterus |
| 细胞定位 | Cytoplasm,Endoplasmic reticulum,Nucleus |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|---|
| 抗原信息 | Recombinant fusion protein corresponding to Human CRABP2. |
| 序列 | VLGVNVMLRKIAVAASKPAVEIKQEGDTFYIKTSTTVRTEINFKVGEEFEEQTVDGRPCKSLVKWESENKM |

靶点信息

| | |
|------|---|
| 研究背景 | This gene encodes a member of the retinoic acid (RA, a form of vitamin A) binding protein family and lipocalin/cytosolic fatty-acid binding protein family. The protein is a cytosol-to-nuclear shuttling protein, which facilitates RA binding to its cognate receptor complex and transfer to the nucleus. It is involved in the retinoid signaling pathway, and is associated with increased circulating low-density lipoprotein cholesterol. Alternatively spliced transcript variants encoding the same protein have been found for this gene. |
|------|---|

| | |
|-------|----------------------|
| 基因ID | 1382 |
| 基因名 | CRABP2 |
| Swiss | P29373 |
| 别名 | CRABP2;CRABP-II;RBP6 |

产品验证



Western blot analysis of CRABP2 expressed in Mouse brain, MCF7, SW480 using CRABP2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn