

— ABLYBIO, Help Your Research



Hsp70 Rabbit mAb

货号: **AYM28729**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	70kDa
实测分子量	70kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,HepG2,BxPC-3,293T,A-549,Mouse brain,Mouse kidney,Mouse lung,Rat lung
细胞定位	blood microparticle,centriole,centrosome,cytoplasm,cytosol,endoplasmic reticulum,extracellular exosome,extracellular region,focal adhesion,mitochondrion,nuclear speck,nucleoplasm,nucleus,perinuclear region of cytoplasm,plasma membrane
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Hsp70 .
------	---


靶点信息

研究背景	This intronless gene encodes a 70kDa heat shock protein which is a member of the heat shock protein 70 family. In conjunction with other heat shock proteins, this protein stabilizes existing proteins against aggregation and mediates the folding of newly translated proteins in the cytosol and in organelles. It is also involved in the ubiquitin-proteasome pathway through interaction with the AU-rich element RNA-binding protein 1. The gene is located in the major histocompatibility complex class III region, in a cluster with two closely related genes which encode similar proteins.
基因ID	3303,3304
基因名	HSPA1A
Swiss	P0DMV8
别名	Hsp70,Hsp70 Rabbit mAb,HSPA1A,Heat shock 70 kDa protein 1,Heat shock protein family A member 1A,HSP72,HSPA1,HSX70


产品验证




Western blot analysis of Hsp70 expressed in Mouse brain,Rat lung,HeLa using Hsp70 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.




Immunohistochemical analysis of paraffin-embedded Human squamous carcinoma, using the Antibody at 1:200 dilution.



Immunohistochemical analysis of paraffin-embedded Human testis cancer, using the Antibody at 1:200 dilution.



Immunohistochemical analysis of paraffin-embedded Human prostate cancer, using the Antibody at 1:200 dilution.



Immunohistochemical analysis of paraffin-embedded human breast, using Hsp70 Antibody .

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)