

# EBI3 Rabbit mAb

货号: **AYM28732**

## 产品信息

|       |  |
|-------|--|
| 反应    | Human,Mouse,Rat  |
| 宿主    | Rabbit   |
| 克隆性   | Monoclonal   |
| 预测反应  |  |
| 应用    | WB   |
| 推荐浓度  | <b>WB:</b> 1:500 - 1:2000  |
| 理论分子量 | 25kDa  |
| 实测分子量 | 25kDa  |
| 形式    | Liquid   |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.                          |
| 偶联物   | Unconjugated   |
| 阳性对照  | MCF7,Mouse spleen,Rat brain  |
| 细胞定位  | endoplasmic reticulum lumen,external side of plasma membrane,extracellular region,extracellular space, plasma membrane |
| 纯化    | Affinity purification  |

## 抗原信息


|      |  |
|------|--|
| 抗原信息 | Recombinant fusion protein corresponding to Human EBI3.  |
| 序列   | DPPEGVRLSPLAERQLQVQWEPGSGWPFPEIFSLKYWIRYKRQGAARFHRVGPIEATSFILRAVRPRARYVQVAAQDLTD YGELSDWSLPATATMSLGK |

## 靶点信息

|      |  |
|------|--|
| 研究背景 | This gene was identified by its induced expression in B lymphocytes in response Epstein-Barr virus infection. It encodes a secreted glycoprotein belonging to the hematopoietin receptor family, and heterodimerizes with a 28 kDa protein to form interleukin 27 (IL-27). IL-27 regulates T cell and inflammatory responses, in part by activating the Jak/STAT pathway of CD4+ T cells. [provided by RefSeq, Sep 2008] |
| 基因ID | 10148  |

|       |                      |
|-------|----------------------|
| 基因名   | EBI3                 |
| Swiss | Q14213               |
| 别名    | IL-27B; IL27B; IL35B |

## 产品验证



Western blot analysis of EBI3 expressed in MCF7, Mouse spleen, Rat brain using EBI3 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)