

CD62P Rabbit mAb

货号: **AYM28733**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	90kDa
实测分子量	105,140kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,LO2,Rat liver,Rat brain
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human CD62P.
序列	ISELTNQKEVAAWTYHYSTKAYSWNISRKYCQNRYTDLVAIQNKNEIDYLNKVLPPYSSYIWIGIRKNNKTWTWVGTKKAL TNEAENWADNEPNNKRNNEDCVEIYIKSPSAPGKWNDHCLKKKHALCYTASCQDMSCSKQGECLLETIGNYTCSCYPGFY GPECEYVREC

靶点信息

研究背景	This gene encodes a 140 kDa protein that is stored in the alpha-granules of platelets and Weibel-Palade bodies of endothelial cells. This protein redistributes to the plasma membrane during platelet activation and degranulation and mediates the interaction of activated endothelial cells or platelets with leukocytes. The membrane protein is a calcium-dependent receptor that binds to sialylated forms of Lewis blood group carbohydrate antigens on neutrophils and monocytes. Alternative splice variants may occur but are not well documented.
------	---

基因ID	6403
基因名	SELP
Swiss	P16109
别名	SELP;CD62;CD62P;GMP140;GRMP;LECAM3;PADGEM;PSEL

产品验证



Western blot analysis of CD62P expressed in Jurkat, LO2, Rat liver using CD62P Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn