

# VEGF Receptor 2 Rabbit mAb

货号: **AYM28752**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IP
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IP:</b> 1:20 - 1:50
理论分子量	75kDa/79kDa/151kDa
实测分子量	210,230 kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse heart
细胞定位	Cell junction,Cell membrane,Cytoplasm,Cytoplasmic vesicle,Early endosome,Endoplasmic reticulum,Nucleus,Secreted,Single-pass type I membrane protein
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human VEGF Receptor 2.
序列	RPTFSELVEHLGNLLQANAQQDGKDYIVLPISSETLSMEEDSGLSLPTSPVSCMEEEEVCDPKFHYDNTAGISQYLQNSKRKSRPVSVKTFEDIPLLEEPEVK

## 靶点信息

研究背景	Vascular endothelial growth factor (VEGF) is a major growth factor for endothelial cells. This gene encode s one of the two receptors of the VEGF. This receptor, known as kinase insert domain receptor, is a type III receptor tyrosine kinase. It functions as the main mediator of VEGF-induced endothelial proliferation, survival, migration, tubular morphogenesis and sprouting. The signalling and trafficking of this receptor are regulated by multiple factors, including Rab GTPase, P2Y purine nucleotide receptor, integrin alphaVbeta3, T-cell protein tyrosine phosphatase, etc.. Mutations of this gene are implicated in infantile capillary hemangiomas.
基因ID	3791
基因名	KDR
Swiss	P35968
别名	CD309;FLK1;VEGFR;VEGFR2;VEGF Receptor 2;KDR

## 产品验证



Western blot analysis of VEGF Receptor 2 expressed in Mouse heart using VEGF Receptor 2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)