

Brachyury Rabbit mAb

货号: **AYM28757**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	41kDa/47kDa
实测分子量	47,41kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	H460
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Brachyury.
序列	HKEMMEEPGDSQQPGYSQWGWLLPGTSTLCPANPHPQGGALSLPSTHSCDRYPTLRSHRSPYPSPYAHRNNSPTYSDNSPACLQSHDNWSSLGMPAHPMSMLPVSHNASPPTSSSQYPSLWSVSNNGAVTPGSQAAAVSNGLGAQFFRGSPAHTPLTHPVSAPSSSGSPLYEGAAAATDIVDSQYDAAAQGRLIASWTPVSPPSM

靶点信息

研究背景	The protein encoded by this gene is an embryonic nuclear transcription factor that binds to a specific DN A element, the palindromic T-site. It binds through a region in its N-terminus, called the T-box, and effects transcription of genes required for mesoderm formation and differentiation. The protein is localized to not ochord-derived cells. Two transcript variants encoding different isoforms have been found for this gene.
------	--

基因ID	6862
基因名	T
Swiss	O15178
别名	T;SAVA;TFT

产品验证



Western blot analysis of Brachyury expressed in H460 using Brachyury Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn