

# DR5 Rabbit mAb

货号: **AYM28758**

## 产品信息

|       |   |
|-------|---|
| 反应    | Human   |
| 宿主    | Rabbit  |
| 克隆性   | Monoclonal  |
| 预测反应  |   |
| 应用    | WB IP FC  |
| 推荐浓度  | <b>WB:</b> 1:500 - 1:2000<br><b>IP:</b> 1:20 - 1:50<br><b>FC:</b> 1:20 - 1:50                 |
| 理论分子量 | 12kDa/45kDa/47kDa   |
| 实测分子量 | 40,48kDa  |
| 形式    | Liquid  |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物   | Unconjugated  |
| 阳性对照  | Jurkat  |
| 细胞定位  | Membrane,Single-pass type I membrane protein  |
| 纯化    | Affinity purification   |

## 抗原信息


|      |  |
|------|--|
| 抗原信息 | Recombinant fusion protein corresponding to Human DR5. |
|------|--|

## 靶点信息

|      |   |
|------|---|
| 研究背景 | The protein encoded by this gene is a member of the TNF-receptor superfamily, and contains an intracellular death domain. This receptor can be activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL/APO-2L), and transduces an apoptosis signal. Studies with FADD-deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein. Two transcript variants encoding different isoforms and one non-coding transcript have been found for this gene. |
| 基因ID | 8795  |

|       |           |
|-------|-----------|
| 基因名   | TNFRSF10B |
| Swiss | O14763    |
| 别名    | DR5       |

## 产品验证



Western blot analysis of DR5 expressed in Jurkat using DR5 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)