

# CRKII Rabbit mAb

货号: **AYM28782**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC IP FC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:200 <b>IP:</b> 1:20 - 1:50 <b>FC:</b> 1:20 - 1:50
理论分子量	34kDa
实测分子量	38kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NIH/3T3,C6
细胞定位	Cell membrane,Cytoplasm
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human CRKII.
------	--

## 靶点信息

研究背景	This gene encodes a member of an adapter protein family that binds to several tyrosine-phosphorylated p roteins. The product of this gene has several SH2 and SH3 domains (src-homology domains) and is involv ed in several signaling pathways, recruiting cytoplasmic proteins in the vicinity of tyrosine kinase through SH2-phosphotyrosine interaction. The N-terminal SH2 domain of this protein functions as a positive regul ator of transformation whereas the C-terminal SH3 domain functions as a negative regulator of transform ation. Two alternative transcripts encoding different isoforms with distinct biological activity have been d escribed. [provided by RefSeq, Jul 2008]
基因ID	1398

基因名	CRK
Swiss	P46108
别名	CRKII

## 产品验证



Western blot analysis of CRKII expressed in NIH/3T3,C6 using CRKII Rabbit mAb at 1:1000. Secondary anti body: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% no n-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)