

# Phospho-p95/NBS1 (S343) Rabbit mAb

货号: **AYM28787**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB ICC IP
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>ICC:</b> 1:50 - 1:200 <b>IP:</b> 1:20 - 1:50
理论分子量	84kDa
实测分子量	95kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,Mouse testis,Rat testis
细胞定位	Chromosome,Nucleus,PML body,telomere
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Phospho-p95/NBS1 (S343) .
------	---

## 靶点信息

研究背景	Mutations in this gene are associated with Nijmegen breakage syndrome, an autosomal recessive chromosomal instability syndrome characterized by microcephaly, growth retardation, immunodeficiency, and cancer predisposition. The encoded protein is a member of the MRE11/RAD50 double-strand break repair complex which consists of 5 proteins. This gene product is thought to be involved in DNA double-strand break repair and DNA damage-induced checkpoint activation.
基因ID	4683
基因名	NBN

Swiss	O60934
别名	Phospho-p95/NBS1 (S343)

## 产品验证



Western blot analysis of Phospho-p95/NBS1 (S343) expressed in 293T, Mouse testis, Rat testis using Phospho-p95/NBS1 (S343) Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBS T. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)