

EDD Rabbit mAb

货号: **AYM28790**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	309kDa
实测分子量	309kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,293T,Jurkat,HeLa
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human EDD.
序列	SISAGIPKVGVLMEVWNMNDSCRFQLRSPESLKNMEKASKTTEAKPESKQEPVKTEMGPPSPASTCSDASSIASSASM PYKRRRSTPAPKEEEKVNEEQWSLREVVVEDVKNPVVGKVLKVDGAYVAVKFPGTSSNTNCQNSSGPDADPSSLLQDC RLLRIDELQVVKTTGGTPKVPDCFQRTPKKLCIPEKTEILAVNVDSKGVHAVL

靶点信息

研究背景	This gene encodes a progestin-induced protein, which belongs to the HECT (homology to E6-AP carboxyl terminus) family. The HECT family proteins function as E3 ubiquitin-protein ligases, targeting specific proteins for ubiquitin-mediated proteolysis. This gene is localized to chromosome 8q22 which is disrupted in a variety of cancers. This gene potentially has a role in regulation of cell proliferation or differentiation.
基因ID	51366

基因名	UBR5
Swiss	O95071
别名	UBR5;DD5;EDD;EDD1;HYD

产品验证



Western blot analysis of EDD expressed in U-87MG,293T,Jurkat using EDD Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn