

CD79a Rabbit mAb

货号: **AYM28811**

产品信息

| | |
|-------|--|
| 反应 | Human |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IHC IF/ICC FC |
| 推荐浓度 | WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50 |
| 理论分子量 | 45-55kDa |
| 实测分子量 | 45-55kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | Raji, Daudi, Mouse spleen, Rat spleen, Rat thymus |
| 细胞定位 | Cell membrane, Single-pass type I membrane protein |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|---|
| 抗原信息 | Recombinant fusion protein corresponding to Human CD79a. |
| 序列 | FRKRWQNEKLGLDAGDEYEDENLYEGLNLDCCSMYEDISRGLQGTYQDVGSLNIGDVQLEK |

靶点信息

| | |
|------|--|
| 研究背景 | The B lymphocyte antigen receptor is a multimeric complex that includes the antigen-specific component, surface immunoglobulin (Ig). Surface Ig non-covalently associates with two other proteins, Ig-alpha and Ig-beta, which are necessary for expression and function of the B-cell antigen receptor. This gene encodes the Ig-alpha protein of the B-cell antigen component. Alternatively spliced transcript variants encoding different isoforms have been described. [provided by RefSeq, Jul 2008] |
|------|--|

| | |
|-------|-------------------------------------|
| 基因ID | 973 |
| 基因名 | CD79A |
| Swiss | P11912 |
| 别名 | CD79A;IGA;MB-1;CD79a molecule;CD79a |

产品验证



Western blot analysis of CD79a expressed in Raji,Daudi,Mouse spleen using CD79a Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn