

ORP1 Rabbit mAb

货号: **AYM28814**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	50,108kDa
实测分子量	50,108kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,293T,SH-SY5Y,Mouse brain,Mouse spleen,Mouse heart,Rat liver,Rat brain
细胞定位	cytosol,endosome,extracellular exosome,late endosome
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human ORP1.
序列	SFAMVLNEVDKDMESVIPKTDCLRPDIRAMENGEIDQASEEKKRLEEKQRAARKNRSKSEEDWKTRWFHQGNPYNGA QDWIYSGSYWDRNYFNLPDIY

靶点信息

研究背景	This gene encodes a member of the oxysterol-binding protein (OSBP) family, a group of intracellular lipid receptors. Most members contain an N-terminal pleckstrin homology domain and a highly conserved C-terminal OSBP-like sterol-binding domain, although some members contain only the sterol-binding domain. Transcript variants derived from alternative promoter usage and/or alternative splicing exist; they encode different isoforms. [provided by RefSeq, Jul 2008]
------	---

基因ID	114876
基因名	OSBPL1A
Swiss	Q9BXW6
别名	

产品验证



Western blot analysis of ORP1 expressed in Mouse brain,Rat liver,HeLa using ORP1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn