

— ABLYBIO, Help Your Research



# Phospho-Histone H3 (T3) Rabbit mAb

货号: **AYM28817**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC FC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>ICC:</b> 1:50 - 1:200 <b>FC:</b> 1:20 - 1:50
理论分子量	15kDa
实测分子量	16kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,HT-29,NCI-H460,HeLa,Mouse stomach,Mouse thymus,Rat stomach
细胞定位	Chromosome,Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Phospho-Histone H3 (T3).
------	--


## 靶点信息

研究背景	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene contains introns and its mRNA is polyadenylated, unlike most histone genes. The protein encoded is a replication-independent member of the histone H3 family.
基因ID	3020
基因名	H3-3A, H3-3B
Swiss	P84243
别名	Phospho-Histone H3 (T3), Phospho-Histone H3 (T3) Rabbit mAb, H3-3A, H3-3B, H3.3A, H3F3, H3F3A, H3.3B, H3F3B

## 产品验证



Western blot analysis of Phospho-Histone H3 (T3) expressed in U-87MG, HT-29, NCI-H460 using Phospho-Histone H3 (T3) Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



Immunohistochemical analysis of paraffin-embedded human tonsil, using Phospho-Histone H3 (T3) Antibody.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))