

— ABLYBIO, Help Your Research



Thyroid Peroxidase Rabbit mAb

货号: **AYM28818**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	103kDa
实测分子量	103kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	cell surface,extracellular space,plasma membrane
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Thyroid Peroxidase.
------	---

靶点信息

研究背景	This gene encodes a membrane-bound glycoprotein. The encoded protein acts as an enzyme and plays a central role in thyroid gland function. The protein functions in the iodination of tyrosine residues in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and triiodothyronine. Mutations in this gene are associated with several disorders of thyroid hormonogenesis, including congenital hypothyroidism, congenital goiter, and thyroid hormone organification defect IIA. Multiple transcript variants encoding distinct isoforms have been identified for this gene, but the full-length nature of some variants has not been determined. [provided by RefSeq, May 2011]
基因ID	7173
基因名	TPO
Swiss	P07202 (https://www.uniprot.org/uniprotkb/P07202/entry)
别名	Thyroid Peroxidase,Thyroid Peroxidase Rabbit mAb,TPO

产品验证



Western blot analysis of Thyroid Peroxidase expressed in HUVEC using Thyroid Peroxidase Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)