— ABLYBIO, Help Your Research

NMI Rabbit mAb

货号: AYM28833

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP
推荐浓度	**WB:** 1:500 - 1:2000 **IHC:** 1:50 - 1:200 **IF/ICC:** 1:50 - 1:200 **IP:** 1:20 - 1:50
理论分子量	35kDa
实测分子量	38kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,A-431,K-562
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human NMI.

靶点信息

研究背景	NMYC interactor (NMI) encodes a protein that interacts with NMYC and CMYC (two members of the oncog ene Myc family), and other transcription factors containing a Zip, HLH, or HLH-Zip motif. The NMI protein also interacts with all STATs except STAT2 and augments STAT-mediated transcription in response to cytok ines IL2 and IFN-gamma. The NMI mRNA has low expression levels in all human fetal and adult tissues tes ted except brain and has high expression in cancer cell line-myeloid leukemias.
基因ID	9111
基因名	NMI
Swiss	Q13287 (https://www.uniprot.org/uniprotkb/Q13287/entry)
别名	NMI,NMI Rabbit mAb,N-myc and STAT interactor

产品验证



Western blot analysis of NMI expressed in HeLa,A-431,K-562 using NMI Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)