

MARK3 Rabbit mAb

货号: **AYM28841**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	
实测分子量	86kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	cytoplasm,cytosol,dendrite,extracellular exosome,plasma membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human MARK3.
序列	MSTRTPPTVNERDTENHTSHGDGRQEVTSRTSRSGARCRNSIASCADEQPHIGNYRLLKTIGKGNFAKVKLARHILTGRE VAIKIIDKTQLNPTSLQKL

靶点信息

研究背景	The protein encoded by this gene is activated by phosphorylation and in turn is involved in the phosphorylation of tau proteins MAP2 and MAP4. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2011]
基因ID	4140

基因名	MARK3
Swiss	P27448
别名	CTAK1;KP78;PAR1A;Par-1a;MARK3

产品验证



Western blot analysis of MARK3 expressed in human breast cancer, mouse liver using MARK3 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn