

— ABLYBIO, Help Your Research



PDI Rabbit mAb

货号: **AYM28854**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB FC
推荐浓度	WB: 1:500 - 1:2000 FC: 1:20 - 1:50
理论分子量	57kDa/58kDa
实测分子量	58kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,Mouse liver,Rat liver
细胞定位	Endoplasmic reticulum lumen
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human PDI.
------	--------------------------------------------------------

靶点信息

研究背景	This gene encodes a member of the disulfide isomerase (PDI) family of endoplasmic reticulum (ER) proteins that catalyze protein folding and thiol-disulfide interchange reactions. The encoded protein has an N-terminal ER-signal sequence, two catalytically active thioredoxin (TRX) domains, two TRX-like domains and a C-terminal ER-retention sequence. The protein plays a role in the folding of nascent proteins in the endoplasmic reticulum by forming disulfide bonds through its thiol isomerase, oxidase, and reductase activity.
基因ID	64714
基因名	PDIA2
Swiss	Q13087 (https://www.uniprot.org/uniprotkb/Q13087/entry)
别名	PDI,PDI Rabbit mAb,PDIA2,Pancreas-specific protein disulfide isomerase,PDIP

产品验证



Western blot analysis of PDI expressed in U-87MG, Mouse liver, Rat liver using PDI Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)