

— ABLYBIO, Help Your Research



Complement C3 Rabbit mAb

货号: **AYM28859**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IP
推荐浓度	WB: 1:500 - 1:2000 IP: 1:20 - 1:50
理论分子量	187kDa
实测分子量	187kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2
细胞定位	Secreted
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Complement C3.
------	--

靶点信息

研究背景	Complement component C3 plays a central role in the activation of complement system. Its activation is required for both classical and alternative complement activation pathways. The encoded preproprotein is proteolytically processed to generate alpha and beta subunits that form the mature protein, which is then further processed to generate numerous peptide products. The C3a peptide, also known as the C3a anaphylatoxin, modulates inflammation and possesses antimicrobial activity. Mutations in this gene are associated with atypical hemolytic uremic syndrome and age-related macular degeneration in human patients. [provided by RefSeq, Nov 2015]
基因ID	718
基因名	C3
Swiss	P01024 (https://www.uniprot.org/uniprotkb/P01024/entry)
别名	Complement C3, Complement C3 Rabbit mAb, C3, C3 and PZP-like alpha-2-macroglobulin domain-containing protein 1, C3adesArg, Complement C3b-alpha' chain, CPAMD1

产品验证



Western blot analysis of Complement C3 expressed in HepG2 using Complement C3 Rabbit mAb at 1:100. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)