

Phospho-eIF4B (S406) Rabbit mAb

货号: **AYM28877**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	64kDa/69kDa
实测分子量	75kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse kidney,Rat skeletal muscle
细胞定位	cytosol,eukaryotic translation initiation factor 4F complex
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Phospho-eIF4B (S406).
序列	SQSTRAASIFGGAKPVDTAAREREVEERLQKEQEKLRQLDEPKLERRPRERHPSWRSEETQERERSRTGSESSQTGTST TSSRNARRRESEKSLNETLNKEEDCHSPTS KPPKPDQPLKVMPAPPPKENAWVKRSSNPPARSQSSDTEQQSPTSGGG KVAPAQPSEEGPGRKDENKVDGMNAPKGQTGNSSRGPDGGNRDHWKESDRKDGKKDQDSRSAPEPKKPEENPASK FSSASKYAALSVDGEDENEGEDYAE

靶点信息

研究背景	Required for the binding of mRNA to ribosomes. Functions in close association with EIF4-F and EIF4-A. Binds near the 5'-terminal cap of mRNA in presence of EIF-4F and ATP. Promotes the ATPase activity and the ATP-dependent RNA unwinding activity of both EIF4-A and EIF4-F.
基因ID	1975

基因名	EIF4B
Swiss	P23588
别名	EIF4B;EIF-4B;PRO1843

产品验证



Western blot analysis of Phospho-eIF4B (S406) expressed in Mouse kidney,Rat skeletal muscle using Phospho-eIF4B (S406) Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn