

— ABLYBIO, Help Your Research



# AHCY Rabbit mAb

货号: **AYM28890**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	44kDa/47kDa
实测分子量	47kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse liver,Mouse pancreas,Rat liver
细胞定位	Cytoplasm,Melanosome
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human AHCY.
------	---------------------------------------------------------

## 靶点信息

研究背景	S-adenosylhomocysteine hydrolase belongs to the adenosylhomocysteinase family. It catalyzes the reversible hydrolysis of S-adenosylhomocysteine (AdoHcy) to adenosine (Ado) and L-homocysteine (Hcy). Thus, it regulates the intracellular S-adenosylhomocysteine (SAH) concentration thought to be important for transmethylation reactions. Deficiency in this protein is one of the different causes of hypermethioninemia. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
基因ID	191
基因名	AHCY
Swiss	P23526 ( <a href="https://www.uniprot.org/uniprotkb/P23526/entry">https://www.uniprot.org/uniprotkb/P23526/entry</a> )
别名	AHCY,AHCY Rabbit mAb,S-adenosyl-L-homocysteine hydrolase,SAHH

## 产品验证



Western blot analysis of AHCY expressed in Mouse liver, Mouse pancreas, Rat liver using AHCY Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))