

# NCTR2 Rabbit mAb

货号: **AYM28894**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	44kDa
实测分子量	44kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,Raji,THP-1
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human NCTR2.
序列	MAWRALHPLLLLLLFPGSQAQSKAQLVLSVAGQTLTVRCQYPPTGSLYEKKGWCKEASALVCIRLVTSSKPRTMAWTSR FTIWDDPDAGFFVTMTDLR

## 靶点信息

研究背景	Cytotoxicity-activating receptor that may contribute to the increased efficiency of activated natural killer (NK cells to mediate tumor cell lysis.
基因ID	9436
基因名	NCR2

Swiss	O95944
别名	NCR2; CD336; LY95; NK-p44; NKP44; dj149M18.1; natural cytotoxicity triggering receptor 2

## 产品验证



Western blot analysis of NCTR2 expressed in Jurkat,Raji,THP-1 using NCTR2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)