

KCNA1 Rabbit mAb

货号: **AYM28895**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IP
推荐浓度	WB: 1:500 - 1:2000 IP: 1:20 - 1:50
理论分子量	56kDa
实测分子量	56kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Cell junction,Cell membrane,Cell projection,Cytoplasmic vesicle,Endoplasmic reticulum,Membrane,Multi-pass membrane protein,Perikaryon,axon,dendrite,presynaptic cell membrane,synapse
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human KCNA1.
序列	QLLHVSSPNLASDSDLRRSSSTMSKSEYMEIEEDMNNSIAHYRQVNIRTANCTTANQNCVNKSKLLTDV

靶点信息

研究背景	This gene encodes a voltage-gated delayed potassium channel that is phylogenetically related to the <i>Drosophila</i> Shaker channel. The encoded protein has six putative transmembrane segments (S1-S6), and the loop between S5 and S6 forms the pore and contains the conserved selectivity filter motif (GYGD). The functional channel is a homotetramer. The N-terminus of the channel is associated with beta subunits that can modify the inactivation properties of the channel as well as affect expression levels. The C-terminus of the channel is complexed to a PDZ domain protein that is responsible for channel targeting. Mutations in this gene have been associated with myokymia with periodic ataxia (AEMK).
------	--

基因ID	3736
基因名	KCNA1
Swiss	Q09470
别名	KCNA1;AEMK;EA1;HBK1;HUK1;KV1.1;MBK1;MK1;RBK1

产品验证



Western blot analysis of KCNA1 expressed in CACO2 using KCNA1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn