

KAT8 / MYST1 / MOF Rabbit mAb

货号: **AYM28903**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	52kDa/53kDa
实测分子量	52kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Rat lung
细胞定位	Chromosome,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human KAT8 / MYST1 / MOF.
序列	MAAQGAAA AAGTSGVAGEGEPGGENAAAEGTAPSPGRVSPPTPARGEPVTV EIGETYL CRRPDSTWHS AEVIQSR VNDQEGREEFYVHYVGFNRR LDEWVDKNRLALTKTVKDAVQKNSEKYLSELA EQPERKITRNQKRKHDEINH VQKTYAEM DPTTAALEKEHEAITKVYVDKI HIGNYEIDAWYFSPFPED

靶点信息

研究背景	This gene encodes a member of the MYST histone acetylase protein family. The encoded protein has a characteristic MYST domain containing an acetyl-CoA-binding site, a chromodomain typical of proteins which bind histones, and a C2HC-type zinc finger. Multiple transcript variants encoding different isoforms have been found for this gene.
------	---

基因ID	84148
基因名	KAT8
Swiss	Q9H7Z6
别名	KAT8;MOF;MYST1;ZC2HC8;hMOF

产品验证



Western blot analysis of KAT8 / MYST1 / MOF expressed in HeLa,Rat lung using KAT8 / MYST1 / MOF R abbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn