

DPYD Rabbit mAb

货号: **AYM28920**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	18kDa/111kDa
实测分子量	110kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	THP-1,Mouse liver
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human DPYD.
序列	MAPVLSKDSADIESILALNPRTQTHATLCSTSAAKLDKHKHWKRNPKNCFNCEKLENNFDDIKHTTLGERGALREAMRCLK CADAPCQKSCPTNLDIKSFITSIANKNYGAAKMIFSDNPLGLTTCGMVCPTSDLCVGGCNLYATEEGPINIGGLQQFATETLI LAFSLMNHL

靶点信息

研究背景	The protein encoded by this gene is a pyrimidine catabolic enzyme and the initial and rate-limiting factor in the pathway of uracil and thymidine catabolism. Mutations in this gene result in dihydropyrimidine dehydrogenase deficiency, an error in pyrimidine metabolism associated with thymine-uraciluria and an increased risk of toxicity in cancer patients receiving 5-fluorouracil chemotherapy. Two transcript variants encoding different isoforms have been found for this gene.
------	--

基因ID	1806
基因名	DPYD
Swiss	Q12882
别名	DPYD;DHP;DHPDHASE;DPD

产品验证



Western blot analysis of DPYD expressed in THP-1, Mouse liver using DPYD Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn