

Cytokeratin 9 Rabbit mAb

货号: **AYM28929**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	62kDa
实测分子量	62kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain
细胞定位	cytoskeleton,cytosol,extracellular exosome,extracellular space,nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Cytokeratin 9.
序列	TANEKSTMQELNSRLASYLDKVVQALEEANNNDLENKIQDWYDKKGPAAIQKNYSPYYNTIDDLKDQIVDLTVGNKTKLLDIDNTRMTLDDFRIFEMEQLNRQGVADANGLRQVLDNLTMEKSDLEMQYETLQEELMALKKNHKEEMSQLTGQNSGDVNVEINVAPGKDLTKTLNDRQYEYEQLIAKNRKDIENTQYETQITQIEHEVSSSGQEVQSSAKEVTQLRHGVQELEIELQSQLSKKAALEKSLEDTKNRYCGQLQMIQEQISNLEAQITDVRQEIECQNQEYSLLSIKMRLEKE

靶点信息

研究背景	This gene encodes the type I keratin 9, an intermediate filament chain expressed only in the terminally differentiated epidermis of palms and soles. Mutations in this gene cause epidermolytic palmoplantar keratoderma.
------	---

基因ID	3857
基因名	KRT9
Swiss	P35527
别名	KRT9;CK-9;EPPK;K9;keratin 9

产品验证



Western blot analysis of Cytokeratin 9 expressed in Mouse brain using Cytokeratin 9 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn