

# WDR4 Rabbit mAb

货号: **AYM28930**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200 <b>FC:</b> 1:20 - 1:50
理论分子量	29kDa/45kDa
实测分子量	45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,HepG2,Rat lung
细胞定位	Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human WDR4.
序列	SRFLATSIASSDDDSLFIYDCSAAEKKSQENKGEDAPLDQGGGAILASTFSKSGSYFALTDDSKRLILFRTPWQCLSVRTV ARRCTALTFIASEEKVLVADKSGDVYFSVLEPHGCGRLELGHLSMLLDVAVSPDDRFILTADRDEKIRVSWAAAPHSIESF CLGHTEFVSRSISVVPTQPGLLLSSSGDGLRLWEYRSGRQLHCCHLASLQELVDPQAPQKFAASRIA

## 靶点信息

研究背景	This gene encodes a member of the WD repeat protein family. WD repeats are minimally conserved regions of approximately 40 amino acids typically bracketed by gly-his and trp-asp (GH-WD), which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this family are involved in a variety of cellular processes, including cell cycle progression, signal transduction, apoptosis, and gene regulation. This gene is excluded as a candidate for a form of nonsyndromic deafness (DFNB10), but is still a candidate for other disorders mapped to 21q22.3 as well as for the development of Down syndrome phenotypes. Alternatively spliced transcript variants have been found for this gene.
基因ID	10785
基因名	WDR4
Swiss	P57081
别名	WDR4;TRM82;TRMT82

## 产品验证



Western blot analysis of WDR4 expressed in HeLa, HepG2, Rat lung using WDR4 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)