

UCHL3 Rabbit mAb

货号: **AYM28959**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	26kDa
实测分子量	26kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y,HT-29,Mouse skeletal muscle,Mouse testis,Rat testis
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human UCHL3 .
序列	MEGQRWLPLEANPEVTNQFLKQLGLHPNWQFVDVYGMPELLSMVPRPVCAVLLLPITEKEYEVRTEEEKIKSQGDV TSSVYFMKQTISNACGTIGLIHAIANNKDKMHFESGSLKFFLEESVMSPEERARYLENYDAIRVTHETSAHEGQTEAPSID EKVDLHFIALVHVDGHLIELDGRKPPINHGGETSDETLLEDAIEVCKKFMERDPDEL

靶点信息

研究背景	The protein encoded by this gene is a member of the deubiquitinating enzyme family. Members of this family are proteases that catalyze the removal of ubiquitin from polypeptides and are divided into five classes, depending on the mechanism of catalysis. This protein may hydrolyze the ubiquitinyl-N-epsilon amide bond of ubiquitinated proteins to regenerate ubiquitin for another catalytic cycle. Alternative splicing results in multiple transcript variants that encode different protein isoforms.
基因ID	7347

基因名	UCHL3
Swiss	P15374
别名	UCHL3;UCH-L3

产品验证



Western blot analysis of UCHL3 expressed in Mouse skeletal muscle,Rat testis,SH-SY5Y using UCHL3 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30 ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 1 20s.

实验步骤

访问官网浏览详情: www.ablybio.cn