

Ku70 Rabbit mAb

货号: **AYM28963**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	65kDa/69kDa
实测分子量	70kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	COS-7,SW480,Mouse brain
细胞定位	Chromosome,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Ku70.
------	---

靶点信息


研究背景	The p70/p80 autoantigen is a nuclear complex consisting of two subunits with molecular masses of approximately 70 and 80 kDa. The complex functions as a single-stranded DNA-dependent ATP-dependent heli case. The complex may be involved in the repair of nonhomologous DNA ends such as that required for d ouble-strand break repair, transposition, and V(D)J recombination. High levels of autoantibodies to p70 an d p80 have been found in some patients with systemic lupus erythematosus.
基因ID	2547

基因名	XRCC6
Swiss	P12956
别名	Ku70

产品验证



Western blot analysis of Ku70 expressed in COS-7, SW480, Mouse brain using Ku70 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30 ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



Immunohistochemical analysis of paraffin-embedded human tonsil, using Ku70 Antibody.

实验步骤

访问官网浏览详情: www.ablybio.cn